

[Go to Product page](#)

## Datasheet for ABIN5526719

**TBK1 ELISA Kit**

## Overview

Quantity:	96 tests
Target:	TBK1
Binding Specificity:	pSer172, total
Reactivity:	Human, Mouse, Rat
Method Type:	Cell ELISA
Application:	ELISA

## Product Details

Purpose:	Cell-Based ELISA Kit. This assay semi-quantitatively measures TBK1 phosphorylated at Serine-172 as well as total TBK1 in adherent cell lines.
Sample Type:	Adherent Cell Culture
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human, Mouse and Rat TBK1 phosphorylated at site Serine-172 as well as total TBK1.
Characteristics:	<ul style="list-style-type: none"><li>• Rapidly measure phosphorylated protein in adherent cell lines</li><li>• Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)</li><li>• No sample lysis is needed</li><li>• Compatible with a standard ELISA plate reader</li></ul>
Components:	<ul style="list-style-type: none"><li>• Uncoated 96-well Strip Microplate</li></ul>

## Product Details

- Wash Buffers
- Fixing Solution
- Quenching Buffer
- Blocking Buffer
- Anti-phospho antibody
- Anti-pan antibody
- HRP-Conjugated Secondary Antibody
- TMB One-Step Substrate
- Stop Solution

### Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

## Target Details

Target: TBK1

Alternative Name: TBK1 ([TBK1 Products](#))

Gene ID: 29110

UniProt: [Q9UHD2](#), [Q9WUN2](#), [D4A7D3](#)

Pathways: [TLR Signaling](#), [Activation of Innate immune Response](#), [Hepatitis C](#), [Toll-Like Receptors Cascades](#), [SARS-CoV-2 Protein Interactome](#)

## Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100 µL of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 °C.
4. Add 100 µL of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.

## Application Details

---

7. Incubate 1 h at RT.
8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 µL of Stop Solution to each well.
11. Read at 450 nm immediately.

---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

---

Storage:	-20 °C
----------	--------

Storage Comment:	Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.
------------------	--

---

Expiry Date:	6 months
--------------	----------