



Datasheet for ABIN5526721

PRKAA1 ELISA Kit



[Go to Product page](#)

3 Images

Overview

Quantity:	96 tests
Target:	PRKAA1
Binding Specificity:	pSer487
Reactivity:	Human, Mouse, Rat
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human, Mouse and Rat Phospho-AMPKa1 (Ser487) ELISA Kit. This assay semi-quantitatively measures AMPKa1 phosphorylated at Serine-487 in cell lysate samples.
Sample Type:	Cell Culture Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human, Mouse and Rat AMPKa1 phosphorylated at site Serine-487.
Characteristics:	<ul style="list-style-type: none">• Pre-Coated 96-well Strip Microplate• Wash Buffer• Anti-Phospho Antibody• HRP-Conjugated Secondary Antibody• Assay Diluent• TMB One-Step Substrate• Stop Solution• Lysis Buffer

Product Details

- Positive Control Sample

- Components:
- Pre-Coated 96-well Strip Microplate
 - Wash Buffer
 - Anti-Phospho Antibody
 - HRP-Conjugated Secondary Antibody
 - Assay Diluent
 - TMB One-Step Substrate
 - Stop Solution
 - Lysis Buffer
 - Positive Control Sample

- Material not included:
- Distilled or deionized water
 - 100 mL and 1 liter graduated cylinders
 - Tubes to prepare sample dilutions
 - Protease and Phosphatase inhibitors
 - Precision pipettes to deliver 2 μ L to 1 mL volumes
 - Adjustable 1-25 mL pipettes for reagent preparation
 - Benchtop rocker or shaker
 - Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target: PRKAA1

Alternative Name: AMPKa1 ([PRKAA1 Products](#))

Gene ID: 5562

UniProt: [Q13131](#), [Q5EG47](#), [P54645](#)

Pathways: [AMPK Signaling](#), [Carbohydrate Homeostasis](#), [Regulation of Carbohydrate Metabolic Process](#), [Warburg Effect](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100 μ L of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 $^{\circ}$ C.
4. Add 100 μ L of prepared primary antibody to each well.

Application Details

5. Incubate 1 h at RT.
6. Add 100 μ L of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.
8. Add 100 μ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 μ L of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only


Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.


Expiry Date: 6 months

Images




ELISA

Image 1. HeLa cells were treated with Calyculin A. Solubilize cells at 4×10^7 cells/ml in Cell Lysate Buffer. Serial dilutions of lysates were analyzed in this ELISA.



ELISA

Image 2. HeLa cells were treated or untreated with Calyculin A. Cell lysates were analyzed using this phosphoELISA and Western Blot.



ELISA

Image 3. HeLa cells were treated or untreated with Calyculin A. Cell lysates were analyzed using this phosphoELISA and Western Blot.