



[Go to Product page](#)

## Datasheet for ABIN5526729

### MARCKS ELISA Kit

#### Overview

Quantity:	96 tests
Target:	MARCKS
Binding Specificity:	pSer152, pSer156, total
Reactivity:	Human, Mouse
Method Type:	Cell ELISA
Application:	ELISA

#### Product Details

Purpose:	Cell-Based ELISA Kit. This assay semi-quantitatively measures MARCKS phosphorylated at Serine-152/156 as well as total MARCKS in adherent cell lines.
Sample Type:	Adherent Cell Culture
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human and Mouse MARCKS phosphorylated at site Serine-152/156 as well as total MARCKS.
Characteristics:	<ul style="list-style-type: none"> <li>• Rapidly measure phosphorylated protein in adherent cell lines</li> <li>• Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)</li> <li>• No sample lysis is needed</li> <li>• Compatible with a standard ELISA plate reader</li> </ul>
Components:	<ul style="list-style-type: none"> <li>• Uncoated 96-well Strip Microplate</li> </ul>

## Product Details

- Wash Buffers
- Fixing Solution
- Quenching Buffer
- Blocking Buffer
- Anti-phospho antibody
- Anti-pan antibody
- HRP-Conjugated Secondary Antibody
- TMB One-Step Substrate
- Stop Solution

Material not included:	<ul style="list-style-type: none"><li>• Distilled or deionized water</li><li>• 100 mL and 1 liter graduated cylinders</li><li>• Tubes to prepare sample dilutions</li><li>• Protease and Phosphatase inhibitors</li><li>• Precision pipettes to deliver 2 µL to 1 mL volumes</li><li>• Adjustable 1-25 mL pipettes for reagent preparation</li><li>• Benchtop rocker or shaker</li><li>• Microplate reader capable of measuring absorbance at 450 nm</li></ul>
------------------------	--

## Target Details

Target:	MARCKS
Alternative Name:	MARCKS ( <a href="#">MARCKS Products</a> )
Gene ID:	4082
UniProt:	<a href="#">P29966</a> , <a href="#">P26645</a>

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Plate:	Pre-coated
Protocol:	<ol style="list-style-type: none"><li>1. Prepare all reagents and samples as instructed in the manual.</li><li>2. Add 100 µL of sample or positive control to each well.</li><li>3. Incubate 2.5 h at RT or O/N at 4 °C.</li><li>4. Add 100 µL of prepared primary antibody to each well.</li><li>5. Incubate 1 h at RT.</li><li>6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.</li><li>7. Incubate 1 h at RT.</li><li>8. Add 100 µL of TMB One-Step Substrate Reagent to each well.</li><li>9. Incubate 30 min at RT.</li></ol>

Application Details

- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions: For Research Use only

Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date: 6 months