



[Go to Product page](#)

Datasheet for ABIN5526738  
**Caveolin-1 ELISA Kit**

### Overview

Quantity: 96 tests

Target: Caveolin-1 (CAV1)

Binding Specificity: pTyr14, total

Reactivity: Human

Method Type: Cell ELISA

Application: ELISA

### Product Details

Purpose: Human Phospho-NBS1 (Ser343) and Total NBS1 Cell-Based ELISA Cell-Based ELISA Kit. This assay semi-quantitatively measures Caveolin-1 phosphorylated at Tyrosine-14 as well as total Caveolin-1 in adherent cell lines.

Sample Type: Adherent Cell Culture

Analytical Method: Semi-Quantitative

Detection Method: Colorimetric

Specificity: This ELISA kit recognizes Human Caveolin-1 phosphorylated at site Tyrosine-14 as well as total Caveolin-1.

Characteristics:

- Rapidly measure phosphorylated protein in adherent cell lines
- Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)
- No sample lysis is needed
- Compatible with a standard ELISA plate reader

## Product Details

---

Components:	<ul style="list-style-type: none"><li>• Uncoated 96-well Strip Microplate</li><li>• Wash Buffers</li><li>• Fixing Solution</li><li>• Quenching Buffer</li><li>• Blocking Buffer</li><li>• Anti-phospho antibody</li><li>• Anti-pan antibody</li><li>• HRP-Conjugated Secondary Antibody</li><li>• TMB One-Step Substrate</li><li>• Stop Solution</li></ul>
-------------	--

Material not included:	<ul style="list-style-type: none"><li>• Distilled or deionized water</li><li>• 100 mL and 1 liter graduated cylinders</li><li>• Tubes to prepare sample dilutions</li><li>• Protease and Phosphatase inhibitors</li><li>• Precision pipettes to deliver 2 <math>\mu</math>L to 1 mL volumes</li><li>• Adjustable 1-25 mL pipettes for reagent preparation</li><li>• Benchtop rocker or shaker</li><li>• Microplate reader capable of measuring absorbance at 450 nm</li></ul>
------------------------	---

## Target Details

---

Target:	Caveolin-1 (CAV1)
Alternative Name:	Caveolin-1 ( <a href="#">CAV1 Products</a> )
Gene ID:	857
UniProt:	<a href="#">Q03135</a>
Pathways:	<a href="#">Maintenance of Protein Location</a> , <a href="#">Signaling Events mediated by VEGFR1 and VEGFR2</a> , <a href="#">Negative Regulation of Transporter Activity</a> , <a href="#">VEGFR1 Specific Signals</a>

## Application Details

---

Application Notes:	Optimal working dilution should be determined by the investigator.
Plate:	Pre-coated
Protocol:	<ol style="list-style-type: none"><li>1. Prepare all reagents and samples as instructed in the manual.</li><li>2. Add 100 <math>\mu</math>L of sample or positive control to each well.</li><li>3. Incubate 2.5 h at RT or O/N at 4 <math>^{\circ}</math>C.</li><li>4. Add 100 <math>\mu</math>L of prepared primary antibody to each well.</li><li>5. Incubate 1 h at RT.</li></ol>

## Application Details

---

6. Add 100  $\mu$ L of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.
8. Add 100  $\mu$ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50  $\mu$ L of Stop Solution to each well.
11. Read at 450 nm immediately.

---

Restrictions: For Research Use only

## Handling

---

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

---

Expiry Date: 6 months