antibodies -online.com





Datasheet for ABIN5526744

CHEK1 ELISA Kit



\sim			
	N/F	ا/\r14	$\triangle W$

OVEIVIEW	
Quantity:	96 tests
Target:	CHEK1
Binding Specificity:	pSer280, total
Reactivity:	Human
Method Type:	Cell ELISA
Application:	ELISA
Product Details	
Purpose:	Human and Mouse Phospho-Rictor (Thr1135) and Total Rictor Cell-Based ELISA Cell-Based ELISA Kit. This assay semi-quantitatively measures CHK1 phosphorylated at Serine-280 as well as total CHK1 in adherent cell lines.
Sample Type:	Adherent Cell Culture
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human CHK1 phosphorylated at site Serine-280 as well as total CHK1.
Characteristics:	 Rapidly measure phosphorylated protein in adherent cell lines Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose) No sample lysis is needed Compatible with a standard ELISA plate reader

Product Details

Components:

- · Uncoated 96-well Strip Microplate
- · Wash Buffers
- · Fixing Solution
- · Quenching Buffer
- · Blocking Buffer
- · Anti-phospho antibody
- · Anti-pan antibody
- · HRP-Conjugated Secondary Antibody
- · TMB One-Step Substrate
- · Stop Solution

Material not included:

- · Distilled or deionized water
- · 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- · Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- · Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	CHEK1
Alternative Name:	CHK1 (CHEK1 Products)
Gene ID:	1111
UniProt:	014757
Pathways:	p53 Signaling, Apoptosis, Cell Division Cycle, DNA Damage Repair

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.	
Plate:	Pre-coated	
Protocol:	1. Prepare all reagents and samples as instructed in the manual.	
	2. Add 100 µL of sample or positive control to each well.	
	3. Incubate 2.5 h at RT or O/N at 4 °C.	
	4. Add 100 μL of prepared primary antibody to each well.	
	5. Incubate 1 h at RT.	
	6. Add 100 μL of prepared 1X HRP-Streptavidin to each well.	
	7. Incubate 1 h at RT.	

Application Details

Expiry Date:

6 months

8. Add 100 µL of TMB One-Step Substrate Reagent to each well. 9. Incubate 30 min at RT. 10. Add 50 µL of Stop Solution to each well. 11. Read at 450 nm immediately. Restrictions: For Research Use only Handling Storage: -20 °C Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.