

Datasheet for ABIN5530303
anti-ASPSCR1 antibody (N-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	400 µL
Target:	ASPSCR1
Binding Specificity:	AA 65-93, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ASPSCR1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This ASPSCR1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 65-93 amino acids from the N-terminal region of human ASPSCR1.
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	ASPSCR1
Alternative Name:	ASPSCR1 (ASPSCR1 Products)
Background:	<p>This gene is a candidate gene for alveolar soft part sarcoma (ASPS). It has been found that this gene is fused with transcription factor TFE3 gene in ASPS and also in renal cell carcinomas.</p> <p>Several alternatively spliced transcript variants of this gene have been described, but their full</p>

Target Details


	length nature has not been determined.
Molecular Weight:	60 kDa
Gene ID:	79058
UniProt:	Q9BZE9

Application Details

Application Notes:	For WB starting dilution is: 1:1000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.



Western Blotting

Image 1. Western blot analysis in A2058 cell line lysates (35ug/lane).



Western Blotting

Image 2. Western blot analysis of ASPSCR1 using rabbit polyclonal ASPSCR1 Antibody using 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the ASPSCR1 gene.