antibodies .- online.com







anti-FAM96A antibody (C-Term)



Images



()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Alternative Name:

Molecular Weight:

Quantity:	400 μL
Target:	FAM96A
Binding Specificity:	AA 109-138, C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAM96A antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This FA96A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 109-138 amino acids from the C-terminal region of human FA96A.
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	FAM96A

FA96A (FAM96A Products)

18 kDa

Target Details

Gene ID:	84191
UniProt:	Q9H5X1

UniProt:	Q9H5X1	
Application Details		
Application Notes:	For WB starting dilution is: 1:1000	
	For IHC-P starting dilution is: 1:50~100	
	For IF starting dilution is: 1:10~50	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	Supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	


should be handled by trained staff only.

4 °C,-20 °C Storage:

Storage Comment: Store at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care


should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to

prolonged high temperatures.




Western Blotting

Image 1. Western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane).



Western Blotting

Image 2. Western blot analysis in WiDr cell line lysates (35ug/lane).



Immunohistochemistry

Image 3. FA96A Antibody immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.