



[Go to Product page](#)

Datasheet for ABIN5530959  
**anti-CNGA2 antibody (N-Term)**

2 Images

Overview

Quantity:	400 µL
Target:	CNGA2
Binding Specificity:	AA 190-218, N-Term
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CNGA2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This CNGA2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 190-218 amino acids from the N-terminal region of human CNGA2.
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	CNGA2
Alternative Name:	CNGA2 ( <a href="#">CNGA2 Products</a> )
Background:	The protein encoded by this gene represents the alpha subunit of a cyclic nucleotide-gated olfactory channel. The encoded protein contains a carboxy-terminal leucine zipper that mediates channel formation.

## Target Details

---

Molecular Weight: 76 kDa

Gene ID: 1260

UniProt: [Q16280](#)

## Application Details

---

Application Notes: For WB starting dilution is: 1:1000

For IHC-P starting dilution is: 1:10~50

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 0.5 mg/mL


Buffer: Supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C,-20 °C

Storage Comment: Store at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.



### Western Blotting

**Image 1.** Western blot analysis in mouse spleen tissue lysates (35ug/lane).



### Immunohistochemistry

**Image 2.** CNGA2 Antibody immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.