



[Go to Product page](#)

Datasheet for ABIN5532393
anti-ZNF217 antibody (N-Term)

2 Images

Overview

| | |
|----------------------|---------------------------------------|
| Quantity: | 400 µL |
| Target: | ZNF217 |
| Binding Specificity: | AA 92-120, N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ZNF217 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|---------------|---|
| Immunogen: | This ZNF217 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 92-120 amino acids from the N-terminal region of human ZNF217. |
| Isotype: | Ig Fraction |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|--|
| Target: | ZNF217 |
| Alternative Name: | ZNF217 (ZNF217 Products) |
| Background: | ZNF217 binds to the promoters of target genes and functions as repressor. Promotes cell proliferation and antagonizes cell death. Promotes phosphorylation of AKT1 at 'Ser-473'. |

Target Details

Molecular Weight: 115 kDa

Gene ID: 7764

UniProt: [O75362](#)

Application Details

Application Notes: For WB starting dilution is: 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.5 mg/mL

Buffer: Supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Storage Comment: Store at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western Blotting

Image 1. Western Blot at 1:2000 dilution + MCF-7 whole cell lysates Lysates/proteins at 20 ug per lane.



Western Blotting

Image 2. Western blot analysis in T47D cell line lysates (35ug/lane).