

Datasheet for ABIN5533861

anti-CCRL2 antibody (AA 206-234)





Overview

Overview	
Quantity:	400 μL
Target:	CCRL2
Binding Specificity:	AA 206-234
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCRL2 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This CCRL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 206-234 amino acids from the Central region of human CCRL2.
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	CCRL2
Alternative Name:	CCRL2 (CCRL2 Products)
Background:	This gene encodes a chemokine receptor like protein, which is predicted to be a seven
	transmembrane protein and most closely related to CCR1. Chemokines and their receptors
	mediated signal transduction are critical for the recruitment of effector immune cells to the site

Target Details

of inflammation. This gene is expressed at high levels in primary neutrophils and primary
monocytes, and is further upregulated on neutrophil activation and during monocyte to
macrophage differentiation. The function of this gene is unknown. This gene is mapped to the
region where the chemokine receptor gene cluster is located.

Molecular Weight:	40 kDa
Gene ID:	9034

Application Details


000421

Application Notes:	For WB starting dilution is: 1:1000
Restrictions:	For Research Use only

Handling

UniProt:

Format:	Liquid
Concentration:	0.4 mg/mL
Buffer:	Supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.



Western Blotting

Image 1. Western blot analysis in 293 cell line lysates (35ug/lane).