

Datasheet for ABIN5536456

anti-PSMD9 antibody (C-Term)





Go to Product page

_						
	V	\triangle	r۱	/1	\triangle	Λ/
	' V '		ΙV			v v

Overview		
Quantity:	400 μL	
Target:	PSMD9	
Binding Specificity:	AA 176-205, C-Term	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PSMD9 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	This PSMD9 antibody is generated from rabbits immunized with a KLH conjugated synthetic	
	peptide between 176-205 amino acids from the C-terminal region of human PSMD9.	
Isotype:	Ig Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	PSMD9	
Alternative Name:	PSMD9 (PSMD9 Products)	
Background:	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure	
	composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4	
	rings of 28 non-identical subunits, 2 rings are composed of 7 alpha subunits and 2 rings are	

composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator.

Molecular Weight: 25 kDa

Gene ID: 5715

UniProt: 000233

Pathways: Positive Regulation of Peptide Hormone Secretion, Negative Regulation of Hormone Secretion,

Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA, Ubiquitin Proteasome Pathway

should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to

Application Details

Application Notes: For WB starting dilution is: 1:1000

Restrictions: For Research Use only

Handling

Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	Supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care	

prolonged high temperatures.



Western Blotting

Image 1. Western blot analysis in A549 cell line lysates (35ug/lane).