# antibodies - online.com







## anti-HAT1 antibody (C-Term)



Image



$\sim$						
	1//	$\Box$	r۱	/ [	$\triangle$	٨

Quantity:	400 μL
Target:	HAT1
Binding Specificity:	AA 389-419, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HAT1 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	This KAT1 (HAT1) antibody is generated from rabbits immunized with a KLH conjugated
	This KAT1 (HAT1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 389-419 amino acids from the C-terminal region of human KAT1
	synthetic peptide between 389-419 amino acids from the C-terminal region of human KAT1
Immunogen:	synthetic peptide between 389-419 amino acids from the C-terminal region of human KAT1 (HAT1).
Immunogen: Isotype:	synthetic peptide between 389-419 amino acids from the C-terminal region of human KAT1 (HAT1).  Ig Fraction
Immunogen:  Isotype:  Purification:	synthetic peptide between 389-419 amino acids from the C-terminal region of human KAT1 (HAT1).  Ig Fraction  This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
Immunogen: Isotype:	synthetic peptide between 389-419 amino acids from the C-terminal region of human KAT1 (HAT1).  Ig Fraction  This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
Immunogen:  Isotype:  Purification:	synthetic peptide between 389-419 amino acids from the C-terminal region of human KAT1 (HAT1).  Ig Fraction  This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
Immunogen:  Isotype:  Purification:  Target Details	synthetic peptide between 389-419 amino acids from the C-terminal region of human KAT1 (HAT1).  Ig Fraction  This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis

replication-dependent nucleosome assembly. The HAT1 protein contains D, A, and B motifs, which are present in many N-acetyltransferases, including those that acetylate substrates other than histones. The HAT1 holoenzyme consists of 2 subunits: the catalytic 46-kD HAT1 and the accessory p46. The p46 subunit stimulated the activity of HAT1 and bound to core histones. The HAT1 holoenzyme acetylated newly synthesized but not nucleosomal histone H4 at lys5 and lys12, and, to a lesser extent, histone H2A at lys5. HAT1 and p46 polypeptides are located in the nucleus of S-phase cells. HAT1 may play a role in telomeric silencing.


Molecular Weight:	50 kDa
Gene ID:	8520
UniProt:	014929

### **Application Details**

Application Notes:	For WB starting dilution is: 1:1000
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	Supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.



#### **Western Blotting**

Image 1. Western blot analysis of hHAT1-E404 in 293, HepG2, Jurkat cell line lysates (35ug/lane)