

Datasheet for ABIN5539576  
**anti-PKC mu antibody (AA 383-395)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	100 µg
Target:	PKC mu (PRKD1)
Binding Specificity:	AA 383-395
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This PKC mu antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunoprecipitation (IP)

## Product Details

Purpose:	PRKD1 (aa383-395)
Sequence:	QDPDPDHEDA NRT
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

## Target Details

Target:	PKC mu (PRKD1)
---------	----------------

## Target Details


Alternative Name:	PRKD1 ( <a href="#">PRKD1 Products</a> )
Background:	PRKD1, PKC-MU, PKCM, PKD, PRKCM,
Gene ID:	5587
NCBI Accession:	<a href="#">NP_002733</a>
Pathways:	<a href="#">Myometrial Relaxation and Contraction</a>

## Application Details

Application Notes:	Western Blot: In transfected HEK293 transiently expressing Human PKD1 (DYKDDDDK tagged), a band of approx. 100 kDa is observed. The same band is observed using anti- DYKDDDDK antibody and no bands are observed in cells transiently expressing Human PKD2 or Peptide ELISA: antibody detection limit dilution 1:8000.
Comment:	<b>Immunoprecipitation:</b> PKD1 was immunoprecipitated from Hek293T cells (as above). Immunoprecipitates were analyzed by Western Blot probed with the same antibody.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Western Blotting

**Image 1.** HEK293 lysate overexpressing Human DYKDDDDK-tagged PKD1, Human DYKDDDDK-tagged PKD2 or Human GST-tagged PKD3 probed with ABIN5539576 (0.1ug/ml) in top panel, probed with anti-DYKDDDDK in middle panel and probed with anti-GST in bottom panel.



Western Blotting

**Image 2.** HEK293 lysate overexpressing Human DYKDDDDK-tagged PKD1 was used to immunoprecipitate PKD1 with 2ug ABIN5539576. The precipitate was subsequently probed in Western blot using ABIN5539576 at 1ug/ml. The secondary anti-goat picks up the heavy chain of ABIN5539576 used for the immunoprecipitation (annotated as IgG). Data kindly obtained from Dr Peter Storz, Mayo Clinic, USA