# antibodies - online.com







# anti-NNMT antibody (N-Term)





		do to Froduct page

( )	11/	NI	/1/	≥W
	/ // <del> </del>	۰۱ ا	// I E	<b>-</b> ¹ \/ \/

Quantity:	100 μg
Target:	NNMT
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This NNMT antibody is un-conjugated
Application:	Western Blotting (WB), ELISA
Draduat Dataila	

#### **Product Details**

Purpose:	NNMT
Sequence:	TSKDTYLSHF NP
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

#### **Target Details**

|--|

## **Target Details**


Alternative Name:	NNMT (NNMT Products)
Background:	NNMT, nicotinamide N-methyltransferase
Gene ID:	4837, 18113, 300691
NCBI Accession:	NP_006160

## Application Details

Restrictions:	For Research Use only
	Peptide ELISA: antibody detection limit dilution 1:128000.
	MW of 29.6 kDa according to NP_006160.1). Recommended concentration: 0.1-0.3 $\mu$ g/mL.
Application Notes:	Western Blot: Approx 28 kDa band observed in lysates of cell lines A549 and HeLa (calculated

## Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



#### **Western Blotting**

**Image 1.** ABIN5539743 (0.1 $\mu$ g/ml) staining of A549 (A) and HeLA (B) lysates (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.