

Datasheet for ABIN5539746
anti-NOLC1 antibody (Internal Region)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	NOLC1
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This NOLC1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Purpose:	NOLC1
Sequence:	QVNSIKFDSE
Isotype:	IgG
Cross-Reactivity:	Cow, Dog, Human, Mouse, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

Target:	NOLC1
---------	-------

Target Details


Alternative Name:	NOLC1 (NOLC1 Products)
Background:	NOLC1, nucleolar and coiled-body phosphoprotein 1, KIAA0035, NOPP130, NOPP140, NS5ATP13, P130, HCV NS5A trans-regulated protein 13, nucleolar and coiled-body phosphoprotein 1, nucleolar phosphoprotein p130, nucleolar protein p130
Gene ID:	9221, 70769, 64896
NCBI Accession:	NP_004732
Pathways:	Regulation of G-Protein Coupled Receptor Protein Signaling

Application Details

Application Notes:	Immunohistochemistry: In paraffin embedded Human Kidney shows strong nucleolar staining in convoluted tubules. Recommended concentration, 2-4 µg/mL. Peptide ELISA: antibody detection limit dilution 1:16000.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Immunohistochemistry

Image 1. ABIN5539746 (2µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.