antibodies .- online.com







anti-FICD antibody (Internal Region)



()\	ve	r\ /	01	Α .
1 11	$/ \square$	I \/ I	\square	/\/

Overview	
Quantity:	100 μg
Target:	FICD
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This FICD antibody is un-conjugated
Application:	ELISA
Product Details	

Purpose:	HYPE / FICD
Sequence:	PPITIRKEQR SD
Isotype:	IgG
Cross-Reactivity:	Cow, Dog, Human, Mouse, Pig, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Recent

Target Details

Target: FICD

Target Details

Alternative Name:	FICD (FICD Products)
Background:	FICD, FIC domain containing, HIP13, HYPE, MGC5623, UNQ3041, Huntingtin interacting protein E, fic S-phase protein cell division homolog, huntingtin interacting protein 13, huntingtin interactor protein E
Gene ID:	11153, 231630, 288741
NCBI Accession:	NP_009007

Application Details

Application Notes:	Western Blot: Preliminary experiments gave an approx 60 kDa band in lysates of Mouse and Rat
	Liver and of cell line HepG2 after 0.3 μ g/mL antibody staining. Please note that currently we
	cannot find an explanation in the literature for the band we observe
	Peptide ELISA: antibody detection limit dilution 1:16000.

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.