

Datasheet for ABIN5540830

## anti-Isocitrate Dehydrogenase antibody



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	Isocitrate Dehydrogenase (IDH)
Reactivity:	Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Isocitrate Dehydrogenase antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Enzyme Immunoassay (EIA), Dot Blot (DB), Immunodiffusion (ID), Radioimmunoassay (RIA), Immunocytochemistry (ICC)

### Product Details

Immunogen:	Isocitrate Dehydrogenase isolated and purified from Porcine heart. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Specificity:	Isocitrate Dehydrogenase isolated and purified from Porcine heart. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. Species: Pig.
Purification:	Solid Phase Affinity Chromatography

## Target Details

Target:	Isocitrate Dehydrogenase (IDH)
Alternative Name:	isocitrate dehydrogenase,idh ( <a href="#">IDH Products</a> )
Background:	<p>Isocitrate dehydrogenase is an enzyme involved in the citric acid cycle. It is 416 amino acids long with a molecular weight of approximately 45 kDa. Isocitrate dehydrogenase enzymes catalyze the oxidative decarboxylation of isocitrate to produce alfa ketoglutarate. The human genome has 5 IDH genes coding for 3 IDH enzymes. The IDH1 and IDH2 require nicotinamide adeninedinucleotide phosphate (NADP) as co-substrate, whereas IDH3 require nicotinamide adenine dinucleotide (NAD). The IDH2 and 3 are localized in mitochondria and are actively involved in the citric acid cycle (TCA) for energy production in contrast, IDH1 is localized in cytoplasm and peroxisomes where it generates NADPH, reduced form of NADP for biosynthetic and other types of reaction. Since alfa KG and NADPH both are intermediately substrate for a number of cellular process, which allows the possibility of oncogenic or tumor suppressive activities of IDH1.</p>
UniProt:	<a href="#">P20304</a>

## Application Details

Application Notes:	<p>Immunoprecipitation. Can be Used: • In precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques). • To prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of choice. Recommended Working Dilutions: Non-precipitating antibody-binding techniques: 1/1,000-1/30,000.</p>
Restrictions:	For Research Use only

## Handling

Buffer:	PBS, pH 7.2 stabilized with Dextran, without preservatives and foreign proteins
Preservative:	Without preservative
Storage:	4 °C,-20 °C
Storage Comment:	<p>Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing. Shelf life: one year from despatch.</p>
Expiry Date:	12 months