antibodies -online.com





anti-JDP2 antibody (AA 21-120) (Biotin)



Go to Product page

| () | 11/ | IN | /ie | A . |
|-----|-------------------|------|--------|--------|
| | / // | ۱ ات | / (− | ' \/\/ |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | JDP2 | |
| Binding Specificity: | AA 21-120 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This JDP2 antibody is conjugated to Biotin | |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human JDP2 | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | JDP2 |
|-------------------|----------------------|
| Alternative Name: | JDP2 (JDP2 Products) |

Target Details

| Background: | Synonyms: Jun dimerization protein 2, JDP2 | | |
|---------------------|---|--|--|
| | Background: Component of the AP-1 transcription factor that represses transactivation | | |
| | mediated by the Jun family of proteins. Involved in a variety of transcriptional responses | | |
| | associated with AP-1 such as UV-induced apoptosis, cell differentiation, tumorigenesis and | | |
| | antitumogeneris. Can also function as a repressor by recruiting histone deacetylase 3/HDAC3 | | |
| | to the promoter region of JUN. May control transcription via direct regulation of the | | |
| | modification of histones and the assembly of chromatin. | | |
| Gene ID: | 122953 | | |
| UniProt: | Q8WYK2 | | |
| Pathways: | Chromatin Binding | | |
| Application Details | | | |
| Application Notes: | WB 1:300-5000 | | |
| | IHC-P 1:200-400 | | |
| | IHC-F 1:100-500 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C for 12 months. | | |
| Expiry Date: | 12 months | | |