

Datasheet for ABIN5557364

anti-CD31 antibody (AA 28-601) (Biotin)



Go to Product page

| _ | | | |
|-----|-----|----|----|
| () | V/C | rv | ٨/ |
| | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | CD31 (PECAM1) | |
| Binding Specificity: | AA 28-601 | |
| Reactivity: | Human, Mouse | |
| Host: | Mouse | |
| Clonality: | Monoclonal | |
| Conjugate: | This CD31 antibody is conjugated to Biotin | |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) | |
| Product Details | | |
| Immunogen: | Recombinant human CD31 | |
| Clone: | 3B5 | |
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse | |
| Purification: | Purified by Protein G. | |
| Target Details | | |
| Target: | CD31 (PECAM1) | |
| Alternative Name: | CD31 | |
| Background: | Synonyms: Platelet endothelial cell adhesion molecule, PECAM1, PECAM-1, EndoCAM, GPIIA, | |

PECA1, CD31

Background: Induces susceptibility to atherosclerosis (By similarity). Cell adhesion molecule which is required for leukocyte transendothelial migration (TEM) under most inflammatory conditions. Tyr-690 plays a critical role in TEM and is required for efficient trafficking of PECAM1 to and from the lateral border recycling compartment (LBRC) and is also essential for the LBRC membrane to be targeted around migrating leukocytes. Prevents phagocyte ingestion of closely apposed viable cells by transmitting 'detachment' signals, and changes function on apoptosis, promoting tethering of dying cells to phagocytes (the encounter of a viable cell with a phagocyte via the homophilic interaction of PECAM1 on both cell surfaces leads to the viable cell's active repulsion from the phagocyte. During apoptosis, the inside-out signaling of PECAM1 is somehow disabled so that the apoptotic cell does not actively reject the phagocyte anymore. The lack of this repulsion signal together with the interaction of the eat-me signals and their respective receptors causes the attachment of the apoptotic cell to the phagocyte, thus triggering the process of engulfment). Isoform Delta15 is unable to protect against apoptosis. Modulates BDKRB2 activation. Regulates bradykinin- and hyperosmotic shock-induced ERK1/2 activation in human umbilical cord vein cells (HUVEC).

 Gene ID:
 5175

 UniProt:
 P16284

Regulation of Actin Filament Polymerization

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Pathways:

| Format: | Liquid | |
|--------------------|--|--|
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |

Handling

| | handled by trained staff only. |
|------------------|--------------------------------|
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Expiry Date: | 12 months |