

Datasheet for ABIN5557503
anti-ONECUT2 antibody (AA 401-504)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	ONECUT2
Binding Specificity:	AA 401-504
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ONECUT2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin- embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.

Target Details

Target:	ONECUT2
Alternative Name:	ONECUT2 (ONECUT2 Products)

Target Details

Background: Synonyms: One cut domain family member 2, ONECUT2, Hepatocyte nuclear factor 6-beta, HNF-6-beta, One cut homeobox 2, Transcription factor ONECUT-2, OC-2, ONECUT2, HNF6B
Background: Transcriptional activator. Activates the transcription of a number of liver genes such as HNF3B.

Gene ID: 9480

UniProt: [O95948](#)

Application Details

Application Notes: WB 1:300-5000
ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
ICC 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months



Western Blotting

Image 1. Mouse cerebrum lysates probed with ONECUT2 Polyclonal Antibody, Unconjugated at 1:300 overnight at 4°C. Followed by a conjugated secondary antibody at 1:5000 for 90 min at 37°C.

Western Blotting

Image 2. Jurkat lysates probed with ONECUT2 Polyclonal Antibody, Unconjugated at 1:300 overnight at 4°C. Followed by a conjugated secondary antibody at 1:5000 for 90 min at 37°C.