

Datasheet for ABIN5557537
anti-RPS6KA1 antibody (pSer380)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	RPS6KA1
Binding Specificity:	pSer380
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This RPS6KA1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	Synthetic peptide derived from human RSK1(S380), around 350-400aa (phospho S380).
Clone:	7A12
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Purification:	Purified by Protein A.

Target Details

Target:	RPS6KA1
Alternative Name:	RSK1 (RPS6KA1 Products)
Background:	Synonyms: p90 RSK1, p90rsk, p90S6K, pp90RSK1, Ribosomal protein S6 kinase 90kD 1,

Ribosomal protein S6 kinase 90kD polypeptide 1, Ribosomal protein S6 kinase 90 kDa polypeptide 1, Ribosomal protein S6 kinase alpha 1, Ribosomal protein S6 kinase polypeptide 1, Ribosomal S6 kinase 1, RPS6K1 alpha, rps6ka, Rps6ka1, RSK 1, RSK 1 p90, RSK, RSK1, S6K alpha 1, 90 kDa ribosomal protein S6 kinase 1, dJ590P13.1 ribosomal protein S6 kinase, 90kD, polypeptide 1, dJ590P13.1, EC 2.7.11.1, HU 1, HU1, MAP kinase activated protein kinase 1a, MAPKAP kinase 1A, MAPKAPK1A, MGC79981, Mitogen-activated protein kinase-activated protein kinase 1A

Background: Serine/threonine-protein kinase that acts downstream of ERK (MAPK1/ERK2 and MAPK3/ERK1) signaling and mediates mitogenic and stress-induced activation of the transcription factors CREB1, ETV1/ER81 and NR4A1/NUR77, regulates translation through RPS6 and EIF4B phosphorylation, and mediates cellular proliferation, survival, and differentiation by modulating mTOR signaling and repressing pro-apoptotic function of BAD and DAPK1. In fibroblast, is required for EGF-stimulated phosphorylation of CREB1, which results in the subsequent transcriptional activation of several immediate-early genes. In response to mitogenic stimulation (EGF and PMA), phosphorylates and activates NR4A1/NUR77 and ETV1/ER81 transcription factors and the cofactor CREBBP. Upon insulin-derived signal, acts indirectly on the transcription regulation of several genes by phosphorylating GSK3B at 'Ser-9' and inhibiting its activity. Phosphorylates RPS6 in response to serum or EGF via an mTOR-independent mechanism and promotes translation initiation by facilitating assembly of the preinitiation complex. In response to insulin, phosphorylates EIF4B, enhancing EIF4B affinity for the EIF3 complex and stimulating cap-dependent translation. Is involved in the mTOR nutrient-sensing pathway by directly phosphorylating TSC2 at 'Ser-1798', which potently inhibits TSC2 ability to suppress mTOR signaling, and mediates phosphorylation of RPTOR, which regulates mTORC1 activity and may promote rapamycin-sensitive signaling independently of the PI3K/AKT pathway. Mediates cell survival by phosphorylating the pro-apoptotic proteins BAD and DAPK1 and suppressing their pro-apoptotic function. Promotes the survival of hepatic stellate cells by phosphorylating CEBPB in response to the hepatotoxin carbon tetrachloride (CCl4).

Gene ID:	6195
UniProt:	Q15418
Pathways:	MAPK Signaling , Neurotrophin Signaling Pathway , Activation of Innate immune Response , Toll-Like Receptors Cascades


Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 1xTBS (pH 7.4), 1 % BSA, 40 %Glycerol and 0.05 % Sodium Azide.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images



Western Blotting

Image 1. A431 cell lysates probed with RSK1(S380) (7A12) Monoclonal Antibody at 1:1000 overnight at 4°C. Followed by a conjugated secondary antibody.