# antibodies .- online.com





# anti-Influenza A Matrix Protein 2 antibody (Influenza A Virus) (AA 2-60) (Alexa Fluor 750)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	Influenza A Matrix Protein 2 (M2)
Binding Specificity:	AA 2-60
Reactivity:	Influenza A Virus, Virus
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Alexa Fluor 750
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from Influenza A virus H5N1 Matrix Protein-2
Isotype:	IgG
Specificity:	This antibody will recognize many Influenza A virus strains
Cross-Reactivity:	Virus
Cross-Reactivity (Details):	Influenza A virus
Purification:	Purified by Protein A.

#### **Target Details**

Target: Influenza A Matrix Protein 2 (M2)

### **Target Details**

Alternative Name:

Target Type:	Influenza Protein
Background:	Synonyms: Avian influenza Matrix Protein-2, Influenza A virus H7N7 H9N2 H13N6 H16N3 H1N1
	N2N1 H3N2 H2N2
	Background: Forms a proton-selective ion channel that is necessary for the efficient release of
	the viral genome during virus entry. After attaching to the cell surface, the virion enters the cell
	by endocytosis. Acidification of the endosome triggers M2 ion channel activity. The influx of
	protons into virion interior is believed to disrupt interactions between the viral ribonucleoprotein
	(RNP), matrix protein 1 (M1), and lipid bilayers, thereby freeing the viral genome from
	interaction with viral proteins and enabling RNA segments to migrate to the host cell nucleus,
	where influenza virus RNA transcription and replication occur. Also plays a role in viral proteins
	secretory pathway. Elevates the intravesicular pH of normally acidic compartments, such as
	trans-Golgi network, preventing newly formed hemagglutinin from premature switching to the
	fusion-active conformation

Influenza A virus Matrix Protein 2 (M2 Products)

## **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
$\vdash$	land	ling	
1 1	ıaııu	11111	٠

Expiry Date:

12 months