

Datasheet for ABIN5564493

IL-37 Protein (AA 1-218) (His tag)



Overview

Quantity:	10 µg
Target:	IL-37 (IL37)
Protein Characteristics:	AA 1-218
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL-37 protein is labelled with His tag.
Application:	SDS-PAGE (SDS)

Product Details

Purpose:	IL-37 (human) (rec.) (His)
Cross-Reactivity:	Human
Characteristics:	Human IL-37 (aa 1-218) is fused at the C-terminus to a His-tag.
Purity:	>95 % (SDS-PAGE)
Endotoxin Level:	<0.01EU/μg purified protein (LAL test).

Target Details

Target:	IL-37 (IL37)
Alternative Name:	IL-37 (IL37 Products)
Background:	Interleukin-1 Family Member 7, IL-1F7, FIL1 zeta, IL-1X, Interleukin-1 Homolog 4, IL-1H4,

Interleukin-1 zeta, IL-1 zeta, IL-1RP1, IL1F7, FIL1Z, IL1H4, IL1RP1		
IL-37 (IL-1F7, IL-1H4) is an IL-1 family member that is expressed only in certain types of human		
organs and cells such as heart, thymus, testis, kidney, mononuclear cells (PBMCs) and dendritic		
cells. IL-37 is an inhibitor of innate immunity with anti-inflammatory properties. It binds to the		
interleukin-18 receptor (IL-18R) and its co-receptor SIGIRR. IL-37 is secreted as a full-length and		
as a processed form starting from amino acid Val46. It plays an important role as a link		
between innate and adaptive immunity and acts as an inhibitor of autoimmune diseases and		
tumor growth. IL-37 also inhibited Lipopolysaccharide (LPS)-induced immunological reaction		
and LPS-induced osteoclast formation and bone resorption.		

Molecular Weight:

~30kDa (SDS-PAGE)

UniProt:

Q9NZH6

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.1 mg/mL
Buffer:	Contains PBS.
Handling Advice:	After reconstitution, prepare aliquots and store at -20 °C. Avoid freeze/thaw cycles. Centrifuge lyophilized vial before opening and reconstitution. PBS containing at least 0.1 % BSA should be used for further dilutions.
Storage:	4 °C,-20 °C
Storage Comment:	Short Term Storage: +4°C Long Term Storage: -20°C Use & Stability: Stable for at least 6 months after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Expiry Date:	6 months