

Datasheet for ABIN5577887

**anti-FAM98C antibody**[Go to Product page](#)

## 2 Images

## Overview

|              |  |
|--------------|--|
| Quantity:    | 100 µL   |
| Target:      | FAM98C   |
| Reactivity:  | Human  |
| Host:        | Rabbit   |
| Clonality:   | Polyclonal   |
| Conjugate:   | This FAM98C antibody is un-conjugated  |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

## Product Details

|                   |   |
|-------------------|---|
| Purpose:          | Rabbit polyclonal antibody raised against recombinant FAM98C.                           |
| Immunogen:        | Recombinant protein corresponding to amino acids of human FAM98C.                       |
| Sequence:         | EPGAGLRLLR FLCSELQATR LLCLRSLLDP SPRPPLGEGV VEGAGMVQEL DLTQLALGLP<br>RPAPGTPASQ LLQELHA |
| Isotype:          | IgG   |
| Cross-Reactivity: | Human   |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | FAM98C   |
| Alternative Name: | FAM98C ( <a href="#">FAM98C Products</a> )                   |
| Background:       | Full Gene Name: family with sequence similarity 98, member C |

## Target Details

Synonyms: FLJ44669

Gene ID: 147965

## Application Details

Application Notes: Immunohistochemistry (1:200-1:500)  
Western Blot (1:250-1:500)  
The optimal working dilution should be determined by the end user.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: In PBS, pH 7.2 (40 % glycerol, 0.02 % sodium azide)


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C


Storage Comment: Store at 4°C. For long term storage store at -20°C.  
Aliquot to avoid repeated freezing and thawing.

## Images



### Immunohistochemistry

**Image 1.** Immunohistochemical staining of human colon with FAM98C polyclonal antibody shows moderate nuclear positivity in glandular cells at 1:200-1:500 dilution.



Western Blotting

**Image 2.** Western blot analysis of Lane 1: RT-4, Lane 2: U-251 MG, Lane 3: Human Plasma, Lane 4: Liver, Lane 5: Tonsil with FAM98C polyclonal antibody at 1:250-1:500 dilution.