
antibodies - online.com







anti-FSHR antibody (N-Term)



Image



Overview

Quantity:	50 μg
Target:	FSHR
Binding Specificity:	N-Term
Reactivity:	Human, Monkey, Gorilla
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FSHR antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Purpose:	Rabbit polyclonal antibody raised against synthetic peptide of FSHR.
Immunogen:	A synthetic peptide corresponding to 17 amino acid at N-terminus of human FSHR.
Specificity:	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Cross-Reactivity:	Gorilla, Human, Monkey
Cross-Reactivity (Details):	BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Target Details

Target:	FSHR
Alternative Name:	FSHR (FSHR Products)
Background:	Full Gene Name: follicle stimulating hormone receptor


Target Details

	Synonyms: FSHRO,LGR1,MGC141667,MGC141668,ODG1
Gene ID:	2492
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Regulation of Intracellular Steroid Hormone Receptor Signaling, Regulation of Hormone Metabolic Process, Platelet-derived growth Factor Receptor Signaling

Application Details

Application Notes:	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (7-14 µg/mL) The optimal working dilution should be determined by the end user.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	In PBS (0.09 % sodium azide)
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

Images



Immunohistochemistry

Image 1. Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) of human testis tissue with FSHR polyclonal antibody . Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.