# antibodies - online.com







# anti-FZD3 antibody (N-Term)



Image



$\sim$				
	$ V \cap$	r\/I	19	٨

Quantity:	50 μg
Target:	FZD3
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Dog, Pig, Chicken, Hamster, Horse, Monkey, Rabbit, Gorilla
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FZD3 antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Purpose:	Rabbit polyclonal antibody raised against synthetic peptide of FZD3.
Immunogen:	A synthetic peptide corresponding to 18 amino acid at N-terminus of human FZD3.
Specificity:	BLAST analysis of the peptide immunogen showed no homology with other human proteins, except FZD6 (39 %).
Cross-Reactivity:	Chicken, Cow, Dog, Gorilla, Hamster, Horse, Human, Monkey, Mouse, Pig, Rabbit, Rat
Cross-Reactivity (Details):	BLAST analysis of the peptide immunogen showed no homology with other human proteins, except FZD6 (39 $\%$ ).
Target Details	
Target:	FZD3

### **Target Details**

Alternative Name:	FZD3 (FZD3 Products)
Background:	Full Gene Name: frizzled homolog 3 (Drosophila) Synonyms: Fz-3,hFz3
Gene ID:	7976
Pathways:	WNT Signaling, Tube Formation

## Application Details

Application Notes:	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (3-5 μg/mL)	
	The optimal working dilution should be determined by the end user.	
Restrictions:	For Research Use only	

### Handling

Format:	Liquid
Buffer:	In PBS (0.09 % sodium azide)
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Store at 4°C. For long term storage store at -80°C.  Aliquot to avoid repeated freezing and thawing.



#### **Immunohistochemistry**

**Image 1.** Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) of human skin (A) and human skeletal muscle tissue (B) with FZD3 polyclonal antibody. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval.