
antibodies -online.com





anti-GPR68 antibody (2nd Cytoplasmic Domain)



Image



Go to Product page

()	1 /	\sim	rv	11/	11	Α
	1//	⊢	I \/	16	٦,	/\

Overview	
Quantity:	50 µg
Target:	GPR68
Binding Specificity:	2nd Cytoplasmic Domain
Reactivity:	Human, Mouse, Rat, Horse, Rabbit, Monkey, Pig, Hamster, Gorilla
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPR68 antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Product Details	

Purpose:	Rabbit polyclonal antibody raised against synthetic peptide of GPR68.
Immunogen:	A synthetic peptide corresponding to 18 amino acids from 2nd cytoplasmic domain of human GPR68.
Specificity:	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Cross-Reactivity:	Gorilla, Hamster, Horse, Human, Monkey, Mouse, Pig, Rabbit, Rat
Cross-Reactivity (Details):	BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Target Details

Target:	GPR68
Alternative Name:	GPR68 (GPR68 Products)

Target Details

Background:	Full Gene Name: G protein-coupled receptor 68	
	Synonyms: MGC111379,MGC156983,OGR1	
Gene ID:	8111	


Application Details

Application Notes:	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (20 μg/mL)
	The optimal working dilution should be determined by the end user.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	In PBS (0.09 % sodium azide)
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

Images



Immunohistochemistry

Image 1. Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human testis with GPR68 polyclonal antibody . Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.