

Datasheet for ABIN5579260  
**anti-Gpr98 antibody (Extracellular Domain)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	50 µg
Target:	Gpr98
Binding Specificity:	Extracellular Domain
Reactivity:	Human, Gorilla, Horse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Gpr98 antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Western Blotting (WB)

## Product Details

Purpose:	Rabbit polyclonal antibody raised against synthetic peptide of GPR98.
Immunogen:	A synthetic peptide corresponding to 19 amino acids at N-terminal extracellular domain of human GPR98.
Specificity:	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Cross-Reactivity:	Gorilla, Horse, Human
Cross-Reactivity (Details):	BLAST analysis of the peptide immunogen showed no homology with other human proteins.

## Target Details

Target:	Gpr98
Alternative Name:	GPR98 ( <a href="#">Gpr98 Products</a> )

## Target Details

Background:	Full Gene Name: G protein-coupled receptor 98 Synonyms: DKFZp761P0710,FEB4,KIAA0686,MASS1,USH2B,USH2C,VLGR1,VLGR1b
Gene ID:	84059
Pathways:	<a href="#">Sensory Perception of Sound</a>


## Application Details

Application Notes:	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (10-26 µg/mL) The optimal working dilution should be determined by the end user.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	In PBS (0.09 % sodium azide)
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

## Images



### Immunohistochemistry

**Image 1.** Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human kidney, renal tubules with GPR98 polyclonal antibody . Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.