

Datasheet for ABIN5582709  
**anti-LRIG2 antibody**



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	LRIG2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LRIG2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Purpose:	Rabbit polyclonal antibody raised against recombinant LRIG2.
Immunogen:	Recombinant protein corresponding to amino acids of human LRIG2.
Sequence:	IPSYLSSQGT LSEPQEGYSN SEAGSHQQLM PPANGYIHKG TDGGTGTRVI CSDCYDNANI YSRTREYCPY TYIAEEDVLD QTLSSLMVQM PKETYL VHPP QDTTALES LI PSANREPSAF PTNHERISEK KLPSTQMSGE TLQRPVWNIN RELGLPHPPF SQQPVHES PQ LHQNEGLAGR EPDCSASSMS CHRLQDHAF
Isotype:	IgG
Cross-Reactivity:	Human

Target Details

Target:	LRIG2
Alternative Name:	LRIG2 ( <a href="#">LRIG2 Products</a> )

## Target Details

Background:	Full Gene Name: leucine-rich repeats and immunoglobulin-like domains 2 Synonyms: DKFZp451C181,KIAA0806,LIG2
Gene ID:	9860

## Application Details


Application Notes:	Immunohistochemistry (1:10-1:20) Western Blot (1:250-1:500) The optimal working dilution should be determined by the end user.
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling


Format:	Liquid
Buffer:	In PBS, pH 7.2 (40 % glycerol, 0.02 % sodium azide)
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.

## Images



### Western Blotting

**Image 1.** Western blot analysis of Lane 1: RT-4, Lane 2: U-251 MG, Lane 3: Human Plasma, Lane 4: Liver, Lane 5: Tonsil with LRIG2 polyclonal antibody .



### Immunohistochemistry

**Image 2.** Immunohistochemical staining of human skeletal muscle with LRIG2 polyclonal antibody shows strong cytoplasmic positivity in myocytes.