

Datasheet for ABIN5591349

## anti-Zinc Finger Protein 354B (ZNF354B) antibody



[Go to Product page](#)

### 2 Images

#### Overview

Quantity:	100 µL
Target:	Zinc Finger Protein 354B (ZNF354B)
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Western Blotting (WB)

#### Product Details

Purpose:	Rabbit polyclonal antibody raised against recombinant ZNF354B.
Immunogen:	Recombinant protein corresponding to amino acids of human ZNF354B.
Sequence:	CKSTPKMTKS TQTQDSFQEQ IRKRLKRDEP WNFISERSCI YEEKLKKQD KNENLQIISV AHTKILTVDR SHKNVEFGQN FYLKSVFIKQ
Isotype:	IgG
Cross-Reactivity:	Human

#### Target Details

Target:	Zinc Finger Protein 354B (ZNF354B)
Alternative Name:	ZNF354B ( <a href="#">ZNF354B Products</a> )
Background:	Full Gene Name: zinc finger protein 354B Synonyms: FLJ25008,KID2,MGC138316

## Target Details

---

Gene ID: 117608

## Application Details

---

Application Notes: Immunohistochemistry (1:50-1:200)  
Western Blot (1:250-1:500)  
The optimal working dilution should be determined by the end user.

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: In PBS, pH 7.2 (40 % glycerol, 0.02 % sodium azide)

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C. For long term storage store at -20°C.  
Aliquot to avoid repeated freezing and thawing.


## Images

---



### Immunohistochemistry

**Image 1.** Immunohistochemical staining of human colon with ZNF354B polyclonal antibody shows strong nuclear positivity in glandular cells at 1:50-1:200 dilution.



### Western Blotting

**Image 2.** Western blot analysis of Lane 1: Negative control (vector only transfected HEK293T lysate), Lane 2: Over-expression Lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells) with ZNF354B polyclonal antibody at 1:250-1:500 dilution.