



Datasheet for ABIN5596830

## anti-COL3 antibody



[Go to Product page](#)

1 Validation

5 Images

5 Publications

### Overview

Quantity:	100 µg
Target:	COL3
Reactivity:	Human, Cow, Mammalian
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This COL3 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

### Product Details

Immunogen:	Immunogen: Collagen Type III from human and bovine placenta Immunogen Type: Native Protein
Isotype:	IgG
Cross-Reactivity (Details):	Typically negligible cross-reactivity against other types of collagens was detected by ELISA against purified standards. Some class-specific anti-collagens may be specific for three-dimensional epitopes which may result in diminished reactivity with denatured collagen or formalin-fixed, paraffin embedded tissues. This antibody reacts with most mammalian Type III collagens and has negligible cross-reactivity with Type I, II, IV, V or VI collagens. Non-specific cross-reaction of anti-collagen antibodies with other human serum proteins or non-collagen extracellular matrix proteins is negligible.
Purification:	Collagen III Antibody has been prepared by immunoaffinity chromatography using immobilized antigens followed by extensive cross-adsorption against other collagens, human serum proteins and non-collagen extracellular matrix proteins to remove any unwanted specificities.

## Target Details

---

Target:	COL3
Alternative Name:	Collagen Type III ( <a href="#">COL3 Products</a> )
Background:	<p>Synonyms: Collagen type III alpha 1 antibody, Collagen type III alpha antibody, EDS4A antibody, Ehlers Danlos syndrome type IV, autosomal dominant antibody, Fetal collagen antibody, COL3A1</p> <p>Background: Rockland produces highly active antibodies and conjugates to collagens. Collagens are highly conserved throughout evolution and are characterized by an uninterrupted "Glycine-X-Y" triplet repeat that is a necessary part of the triple helical structure. For these reasons, it is often extremely difficult to generate antibodies with specificities to collagens. The development of 'type' specific antibodies is dependent on NON-DENATURED three-dimensional epitopes. Rockland extensively purifies collagens for immunization from human and bovine placenta and cartilage by limited pepsin digestion and selective salt precipitation. This preparation results in a native conformation of the protein. Antibodies are isolated from rabbit antiserum and are extensively cross-adsorbed by immunoaffinity purification to produce 'type' specific antibodies. Greatly diminished reactivity and selectivity of these antibodies will result if denaturing and reducing conditions are used for SDS-PAGE and immunoblotting. Ideal for investigators involved in Cell Biology, Signal Transduction and Stem Cell research.</p> <p>Gene Name: COL3A1</p>
Gene ID:	1281
UniProt:	<a href="#">P02461</a>

## Application Details

---

Application Notes:	<p>Immunohistochemistry Dilution: 1:50 - 1:200</p> <p>Application Note: Anti-Collagen antibodies have been used for indirect trapping ELISA for quantitation of antigen in serum using a standard curve, immunoprecipitation, native (non-denaturing, non-dissociating) PAGE, immunohistochemistry, and western blotting for highly sensitive qualitative analysis.</p> <p>Western Blot Dilution: 1:1,000 - 1:10,000</p> <p>Immunoprecipitation Dilution: 1:100</p> <p>ELISA Dilution: 1:5,000 - 1:50,000</p>
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1.0 mg/mL
Buffer:	Buffer: 0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0 Stabilizer: None
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage, mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Expiry Date:	12 months

## Publications


Product cited in:	Ida, Hikage, Itoh, Ida, Ohguro: "Prostaglandin F2α agonist-induced suppression of 3T3-L1 cell adipogenesis affects spatial formation of extra-cellular matrix." in: <b>Scientific reports</b> , Vol. 10, Issue 1, pp. 7958, (2020) ( <a href="#">PubMed</a> ).
-------------------	--

## Images




### Western Blotting

**Image 1.** Western Blot of Rabbit Anti-COLLAGEN III Antibody Lane 1: Human Collagen III Load: 100 ng per lane  
Primary antibody: Collagen III Antibody at 1:1000 o/n at 4°C  
Secondary antibody: 649 Goat anti-rabbit at 1:20,000 for 30 min at RT  
Block: ABIN925618 for 30 min at RT  
Predicted/Observed size: 138 kDa, 138 kDa



### Immunohistochemistry

**Image 2.** anti collagen III antibody (600-401-105 Lot 26016, 1:400, 45 min RT) showed strong staining in FFPE sections of human skin(left, dermis) with moderate to strong red staining and testis (right) where strong staining was observed within connective tissue between seminiferous tubules. The antibody showed strong extracellular staining within connective tissues across many organs with minimal background staining. Slides were steamed in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes for antigen retrieval. Images provided courtesy of LifeSpan Biosciences, Seattle, WA



### Immunohistochemistry

**Image 3.** Immunohistochemistry of Rabbit Anti-collagen type III antibody. Tissue: right lobe of the liver section. A:Central Vein (CV) fibrosis, B: Non-fibrotic CV, C: Perisinusoidal fibrosis, D: Non-fibrotic area, E: Protoplasmic tract fibrosis, F: Septal fibrosis (arrow). Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: Anti-collagen type III at 1:500 for 4°C for 24hr. Secondary antibody: Peroxidase biotin-streptavidin rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: Anti-collagen type III is intra and extracellular. Staining: 3,3'-diaminobenzidine tetrahydrochloride was used as the chromogen. Nuclei were counterstained purple with hematoxylin.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN5596830.



**Successfully validated (Immunohistochemistry (IHC))**

by [MS Validated Antibodies](#)

Report Number: 300050

Date: Aug 22 2023


Target:	COL3
Lot Number:	47783
Method validated:	Immunohistochemistry (IHC)
Positive Control:	Human TMA Mouse monoclonal anti-COL3 antibody clone FH-7A rabbit monoclonal anti-COL3 antibody HL1906
Notes:	Passed. The anti-COL3 antibody ABIN5596830 stains collagen III in formalin-fixed tissues consistently with expected staining pattern.
Primary Antibody:	ABIN5596830
Secondary Antibody:	EnVision Polymer-HRP mouse/rabbit Kit, Dako REAL, K5007
Protocol:	<ul style="list-style-type: none"><li>• Slide preparation<ul style="list-style-type: none"><li>◦ Mount 2,5 µm FFPE tissue sections on superfrost slides.</li><li>◦ Deparaffinize tissue sections 3x 5 min in xylene.</li><li>◦ Rehydrate tissue sections in a descending ethanol series for 1 min each 100%, 96%, and 80% ethanol.</li><li>◦ Rinse tissue sections for 5 min in TBST buffer (DAKO, K8000).</li></ul></li><li>• Epitope retrieval<ul style="list-style-type: none"><li>◦ Autoclave tissue sections for 5 min at 121 °C in 1x Tris-EDTA-citrate buffer pH7.8 (20x Tris-EDTA-citrate buffer stock solution: 5 g Trizma base (Sigma-Aldrich, T1503), 10 g EDTA (Merck, 1.08418), 6.4g tri-sodium citrate (Sigma-Aldrich, C0909), adjust to pH 7.8 using HCL 1 M, ad 1 L with dH<sub>2</sub>O).</li><li>◦ Rinse tissue sections for 5 min in TBST buffer.</li></ul></li><li>• Peroxidase blocking<ul style="list-style-type: none"><li>◦ Incubate tissue sections for 10 min in Peroxidase-Blocking Solution (Dako REAL, S2023).</li><li>◦ Rinse tissue sections 2x for 5 min in TBST buffer.</li></ul></li><li>• Antibody incubation<ul style="list-style-type: none"><li>◦ Dilute primary rabbit anti-collagen III antibody (antibodies-online, ABIN5596830, lot 47783) diluted 1:112.5, 1:225, or 1:450 in antibody diluent (Dako REAL, S2022).</li><li>◦ Cover tissue section with 100-200 µl diluted antibody.</li><li>◦ Incubate tissue sections for 1 h at 37 °C in a moist chamber.</li></ul></li></ul>

- Rinse tissue sections for 5 min in TBST buffer.
- Apply EnVision Polymer-HRP mouse/rabbit Kit (Dako REAL, K5007) according to manufacturer's recommendation.
- Rinse tissue sections 2x for 5 min in TBST buffer.
- Staining
  - Cover slides for 10 min with DAB-Chromogen (EnVision Polymer-HRP mouse/rabbit Kit, Dako REAL, K5007).
  - Wash slides thoroughly with dH<sub>2</sub>O.
  - Counterstain for 15 sec with Hematoxylin (Mayers Hematoxylin: 200ml ddH<sub>2</sub>O, 0,2g Hematoxylin (Serva, 24420.02), 10 g aluminium potassium sulfate dodecahydrate (Merck, 1.01047), 0,04 g sodium iodate (Merck, 1.06525), 10 g chloral hydrate (Sigma-Aldrich, 15307)).
  - Develop for 15 sec in H<sub>2</sub>O.
  - Dehydrate tissue sections in an ascending ethanol series for 1 min each 80%, 96%, 100% ethanol.
  - Wash tissue sections 3x 5 min in xylene.
  - Apply mounting medium and coverslips.
- Image acquisition
  - Acquire images using a Galileo TMAtic (ISENET).

---


Experimental Notes:

- For antibody comparison an antibody test TMA was used that contained 80 normal tissues from 21 different organs and 95 neoplastic tissues from 18 different tumor types.
  - The anti-COL3 antibody ABIN5596830 tends to cause a rather ubiquitous cytoplasmic staining of all kinds of cells including epithelial cells. This cytoplasmic staining is reduced to a tolerable level at dilutions of 1:225 and less. At such dilutions, ABIN5596830 stains fibrillar structures – also detected by the two reference antibodies - at weak to moderate intensity. Based on the identical staining pattern obtained by three different collagen III antibodies, these fibrillar staining's are likely due to a binding of the antibodies to collagen III. Collagen III staining is typically located adjacent to epithelial structures, in smooth muscles or around vessels of all sizes, and in the stroma of tumors.
  - ABIN5596830 cross-reacts with some inflammatory cells in lymph nodes.
-




**Validation image no. 1 for anti-Collagen, Type III (COL3) antibody (ABIN5596830)**

IHC staining of the stroma in colorectal cancer with anti-collagen III antibody ABIN5596830 diluted 1:225 shows fibrillar collagen III staining.




**Validation image no. 2 for anti-Collagen, Type III (COL3) antibody (ABIN5596830)**

IHC staining of squamous cell carcinoma of the oral cavity with anti-collagen III antibody ABIN5596830 diluted 1:225 shows a distinct fibrillar collagen III staining of the stroma.



**Validation image no. 3 for anti-Collagen, Type III (COL3) antibody (ABIN5596830)**

IHC staining of heart tissue with anti-collagen III antibody ABIN5596830 diluted 1:225 shows distinct fibrillar collagen III staining surrounding each heart muscle cell.



**Validation image no. 4 for anti-Collagen, Type III (COL3) antibody (ABIN5596830)**

IHC staining of non-cancerous breast tissue with anti-collagen III antibody ABIN5596830 diluted 1:225 shows considerable sclerosis.