

Datasheet for ABIN5608232

CD47 Protein (CD47) (partial) (Fc Tag)



_				
()	1//	rv	IO	Λ/
()	VC	. I V	1	v v

Quantity:	100 μg	
Target:	CD47	
Protein Characteristics:	partial	
Origin:	Mouse	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This CD47 protein is labelled with Fc Tag.	
Application:	Western Blotting (WB)	
Product Details		
Sequence:	Gln 19 - Lys 140	
Purity:	>95 % as determined by SDS-PAGE.	
Endotoxin Level:	Endotoxin level is less than 1.0 EU per ug by the LAL method.	
Target Details		
Target:	CD47	
Alternative Name:	e Name: CD47 (CD47 Products)	
Background:	Leukocyte surface antigen CD47 is also known as Antigenic surface determinant protein OA3,	
	Integrin-associated protein (IAP) and Protein MER6. CD47 contains 1 Ig-like V-type	
	(immunoglobulin-like) domain. CD47 is very broadly distributed on normal adult tissues. CD47	

has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in

Target Details

the modulation of integrins and plays an important role in memory formation and synaptic plasticity in the hippocampus by similarity. CD47 is the receptor for SIRPA, binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. CD47 Interaction with SIRPG mediates cell-cell adhesion, enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation.

Molecular Weight: 40 kDa

Gene ID: 16423

UniProt: Q61735

Application Details

Application Notes: This recombinant protein can be used for WB.

Restrictions: For Research Use only

Handling

Format:	Lyophilized	
Buffer:	50 mM Tris, 100 mM Glycine, pH 7.5	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Lyophilized Protein should be stored at -20°C or lower for long term storage. Upon reconstitution, working aliquots should be stored at -20°C or -70°C.	