

Datasheet for ABIN5611444
anti-SKP2 antibody (AA 221-425)



[Go to Product page](#)

3 Images

Overview

| | |
|----------------------|---|
| Quantity: | 0.25 mL |
| Target: | SKP2 |
| Binding Specificity: | AA 221-425 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SKP2 antibody is un-conjugated |
| Application: | Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|---------------|--|
| Immunogen: | SKP2 antibody was raised against recombinant protein encoding aa 221-425 of human Skp2 alpha Genename: SKP2 |
| Specificity: | This antibody reacts to S-phase Kinase-associated Protein 2 (p45) (SKP2). |
| Purification: | Immunoaffinity Chromatography |

Target Details

| | |
|-------------------|---|
| Target: | SKP2 |
| Alternative Name: | SKP2 / FBXL1 (SKP2 Products) |
| Background: | Synonyms: Cyclin-A/CDK2-associated protein p45, F-box/LRR-repeat protein 1, S-phase kinase-associated protein 2 |

Target Details

| | |
|-----------------|--|
| Gene ID: | 6502 |
| NCBI Accession: | NP_005974 |
| UniProt: | Q13309 |
| Pathways: | Mitotic G1-G1/S Phases |


Application Details

| | |
|--------------------|--|
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions: | For Research Use only |

Handling


| | |
|--------------------|--|
| Concentration: | 0,2 mg/mL |
| Buffer: | 10 mM PBS, pH 7.4, BSA, sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |
| Storage Comment: | Store the antibody undiluted at 2-8 °C. DO NOT FREEZE! |

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Human Prostate (formalin-fixed, paraffin-embedded) stained with SKP2at 10 µg/ml followed by biotinylated goat anti-rabbit IgG secondary antibody, alkaline phosphatase-streptavidin and chromogen.



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Human prostate stained with anti-SKP2



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Human Adrenal (formalin-fixed, paraffin-embedded) stained with SKP2at 10 µg/ml followed by biotinylated goat anti-rabbit IgG secondary antibody, alkaline phosphatase-streptavidin and chromogen.