



[Go to Product page](#)

Datasheet for ABIN5650303

## anti-Usp30 antibody

### 2 Images

#### Overview

Quantity:	100 µL
Target:	Usp30
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Usp30 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### Product Details

Purpose:	Mouse monoclonal antibody raised against partial recombinant human USP30.
Immunogen:	Recombinant protein corresponding to human USP30.
Sequence:	IEAKGTLNGE KVEHQRTTFV KQLKLGKLPQ CLCIHLQRLS WSSHGTPLKR HEHVQFNEFL MMDIYKYHLL GHKPSQHNP KLNKNPGPTLE LQDGGAPTP VLNQPGAPKT QIFMNGACSP SLLPTLSAPM PFPLPVVPDY SSST
Clone:	CL4438
Isotype:	IgG2a
Cross-Reactivity:	Human

#### Target Details

Target:	Usp30
---------	-------

## Target Details

Alternative Name:	USP30 ( <a href="#">Usp30 Products</a> )
Background:	Full Gene Name: ubiquitin specific peptidase 30 Synonyms: FLJ40511,MGC10702
Gene ID:	84749


## Application Details

Application Notes:	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:200-1:500) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	In PBS, pH 7.2 (40 % glycerol, 0.02 % sodium azide).
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.

## Images



### Western Blotting

**Image 1.** Western Blot ((Transfected lysate) analysis of (1) control, (2) USP30 over-expression lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells, LY409983).



### Immunohistochemistry

**Image 2.** Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human testis shows moderate cytoplasmic immunoreactivity in seminiferous tubules cells.