



[Go to Product page](#)

Datasheet for ABIN5650750

## anti-GST antibody (Alkaline Phosphatase (AP))

### 1 Image

#### Overview

Quantity:	100 µg
Target:	GST
Reactivity:	Schistosoma japonicum
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This GST antibody is conjugated to Alkaline Phosphatase (AP)
Application:	Western Blotting (WB), ELISA

#### Product Details

Immunogen:	Glutathione-S-Transferase (GST) tagged protein
Clone:	3E2
Isotype:	IgG1
Specificity:	Detects 26 kDa for purified GST protein. Does not detect proteins without a GST tag.
Purification:	Protein G Purified

#### Target Details

Target:	GST
Alternative Name:	GST ( <a href="#">GST Products</a> )
UniProt:	<a href="#">P08515</a>

## Application Details

Application Notes:	<ul style="list-style-type: none"><li>• WB (1:1000)</li><li>• ELISA (1:800)</li><li>• optimal dilutions for assays should be determined by the user.</li></ul>
--------------------	--


Comment:	A 1:1000 dilution of ABIN5650750 was sufficient for detection of GST in 1 µg of Purified GST tagged PmtA by ECL immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.
----------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % Sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C

## Images



### Western Blotting

**Image 1.** Western Blot analysis of *Schistosoma japonicum* Purified GST showing detection of 26 kDa GST protein using Mouse Anti-GST Monoclonal Antibody, Clone 3E2 (ABIN5650750). Lane 1: Molecular Weight Ladder (MW). Lane 2: Purified *Pseudomonas aeruginosa* Metallothionein (PmtA) GST tagged protein. Lane 3: GST. Load: 1 µg. Block: 5 % Skim Milk powder in TBST. Primary Antibody: Mouse Anti-GST Monoclonal Antibody (ABIN5650750) at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG:HRP at 1:5000 for 1 hour at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 26 kDa.