antibodies -online.com





anti-ROR2 antibody (AA 100-200) (FITC)

2 Images



Go to Product page

\sim			
	N/6	1//r	$I \cap V$

Overview	
Quantity:	100 μg
Target:	ROR2
Binding Specificity:	AA 100-200
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ROR2 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	Synthetic peptide of Human ROR2 (100-200 aa), conjugated to Keyhole Limpet Haemocyanin (KLH).
Specificity:	Detects 104.7 kDa. Bands at ~60, 35 kDa are possible degradation products.
Purification:	Peptide Affinity Purified
Target Details	
Target:	ROR2
Alternative Name:	ROR2 (ROR2 Products)
Background:	ROR2 is a receptor tyrosine kinase which may have a role in early cartilage and growth plate development. ROR2 expression is generally highest during embryonic development and will gradually decrease as cells terminally differentiate. Wnt5a has been identified to utilize ROR2 as

ne ID:	4920
	their functional effect has not been well characterized.
	as an oncogene. Primarily mis-sense mutations of ROR2 have been identified in tumours but
	formation was implicated as the mechanism for cancer progression. ROR2 seems to function
	prostate, osteosarcomas and metastatic melanomas, Wnt5a/ROR2 receptor complex
	cell carcinomas. Increased expression of ROR2 was correlated with increased metastasis. In
	implicated in gastric, osteosarcoma, mestatic melanoma, prostate, renal cell and squamous
	a co-receptor for non-canonical Wnt signalling. Aberrant expression of ROR2 has been

Gene ID:	4920
NCBI Accession:	NP_001305133
UniProt:	Q01974
Pathways:	RTK Signaling, WNT Signaling

Application Details


Application Notes:	 WB (1:1000) IHC (1:50) optimal dilutions for assays should be determined by the user.
Comment:	A 1:1000 dilution of ABIN5651092 was sufficient for detection of ROR2 in 10 µg of human HeLa
	cell lysates by ECL immunoblot analysis using goat anti-rabbit IgG:HRP as the secondary

Restrictions: For Research Use only

antibody.


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C



Immunohistochemistry

Image 1. Immunohistochemistry analysis using Rabbit Anti-ROR Polyclonal Antibody (ABIN5651092). Tissue: Prostate. Species: Human. Fixation: Formalin Fixed Paraffin-Embedded. Primary Antibody: Rabbit Anti-ROR Polyclonal Antibody (ABIN5651092) at 1:50 for 30 min at RT. Counterstain: Hematoxylin. Magnification: 20X.



Western Blotting

Image 2. Western blot analysis of Human Cervical cancer cell line (HeLa) lysate showing detection of ~104.7 kDa ROR2 protein using Rabbit Anti-ROR2 Polyclonal Antibody (ABIN5651092). Lane 1: Molecular Weight Ladder (MW). Lane 2: Cervical Cancer cell line (HeLa) lysate. Load: 10 μg. Block: 5 % Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-ROR2 Polyclonal Antibody (ABIN5651092) at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Rabbit HRP:lgG at 1:4000 for 1 hour at RT. Color Development: ECL solution for 5 min at RT. Predicted/Observed Size: ~104.7 kDa. Other Band(s): ~60, 35 kDa degradation products.