

Datasheet for ABIN5651516

Afamin ELISA Kit



Overview

| Quantity: | 96 tests |
|--------------------------|------------------------|
| Target: | Afamin (AFM) |
| Reactivity: | Rat |
| Method Type: | Competition ELISA |
| Detection Range: | 2.47 ng/mL - 200 ng/mL |
| Minimum Detection Limit: | 2.47 ng/mL |
| Application: | ELISA |

Product Details

| Sample Type: | Plasma, Serum |
|--------------------|---|
| Analytical Method: | Quantitative |
| Detection Method: | Colorimetric |
| Specificity: | This assay has high sensitivity and excellent specificity for detection of Afamin (AFM). No significant cross-reactivity or interference between Afamin (AFM) and analogues was observed. |
| Sensitivity: | 0.93 ng/mL |

Target Details

| Target: | Afamin (AFM) |
|-------------------|-----------------------|
| Alternative Name: | Afamin (AFM Products) |
| Background: | Gene Name: Afamin |

Gene Aliases: ALB2, ALBA, ALF, A-ALB, Alpha-Albumin

Application Details

| Application Details | |
|---------------------|--|
| Comment: | The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end. |
| Assay Time: | 2 h |
| Plate: | Pre-coated |
| Protocol: | The microtiter plate provided in this kit has been pre-coated with an antigen. Standards or samples are then added to the appropriate microtiter plate wells with a Horseradish Peroxidase (HRP)-conjugated secondary antibody. After TMB substrate solution is added, those wells that contain Afamin (AFM) will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Afamin (AFM) in the samples is then determined by comparing the O.D. of the samples to the standard curve. |
| Assay Precision: | Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Afamin (AFM) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Afamin (AFM) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12% |
| Restrictions: | For Research Use only |
| Handling | |
| Handling Advice: | The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | -20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at |
| | |

Handling

4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.

Expiry Date:

4-8 months