

Datasheet for ABIN5651558 ALDH1A2 ELISA Kit



Overview

Quantity:	96 tests
Target:	ALDH1A2
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.625 ng/mL - 40 ng/mL
Minimum Detection Limit:	0.625 ng/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2). No significant cross-reactivity or interference between Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2) and analogues was observed.
Sensitivity:	0.215 ng/mL
Target Details	

Target: ALDH1A2 Alternative Name: Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5651558 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Gene Name: Aldehyde Dehydrogenase 1 Family, Member A2
	Gene Aliases: RALDH(II), RALDH2, RALDH2-T, Retinaldehyde Dehydrogenase 2, Retinal
	dehydrogenase 2, Retinaldehyde-specific dehydrogenase type 2
Gene ID:	8854
UniProt:	094788
Pathways:	Retinoic Acid Receptor Signaling Pathway

Application Details

Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2), biotin- conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme- substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2) in the samples is then determined by comparing the 0.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Aldehyde Dehydrogenase 1 Family, Member A2 (ALDH1A2) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5651558 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Application Details	
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months