

Datasheet for ABIN5651675

ALAS1 ELISA Kit



_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	96 tests
Target:	ALAS1
Reactivity:	Rat
Method Type:	Sandwich ELISA
Detection Range:	0.781 ng/mL - 50 ng/mL
Minimum Detection Limit:	0.781 ng/mL
Application:	ELISA

Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Aminolevulinate Delta Synthase 1 (ALAS1). No significant cross-reactivity or interference between Aminolevulinate Delta Synthase 1 (ALAS1) and analogues was observed.
Sensitivity:	0.36 ng/mL

Target Details

Target:	ALAS1	
Alternative Name:	Aminolevulinate Delta Synthase 1 (ALAS1 Products)	

Target Details Gene Name: Aminolevulinate Delta Synthase 1 Background: Gene Aliases: ALAS3, ALAS, ALASH, 5-aminolevulinic acid synthase 1, Delta-ALA synthase 1, 5aminolevulinate synthase, nonspecific, mitochondrial Regulation of Lipid Metabolism by PPARalpha Pathways: **Application Details** Comment: The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end. Assay Time: 3 h Plate: Pre-coated Protocol: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Aminolevulinate Delta Synthase 1 (ALAS1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Aminolevulinate Delta Synthase 1 (ALAS1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Aminolevulinate Delta Synthase 1 (ALAS1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Aminolevulinate Delta Synthase 1 (ALAS1) in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Assay Precision: Aminolevulinate Delta Synthase 1 (ALAS1) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level

Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
Aminolevulinate Delta Synthase 1 (ALAS1) were tested 20 times on one plate, respectively
Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
Aminolevulinate Delta Synthase 1 (ALAS1) were tested on 3 different plates, 8 replicates in each
plate. CV(%) = SD/meanX100
Intra-Assay: CV<10%
Inter-Assay: CV<12%

Restrictions: For Research Use only

Handling

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and	
	specimen samples should be assayed in duplicate. Once the procedure has been started, all	
	steps should be completed without interruption.	
Storage:	4 °C,-20 °C	
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.	
Expiry Date:	4-8 months	