

Datasheet for ABIN5652063

ABCA4 ELISA Kit



_			
()	V/C	rv	٨/

Quantity:	96 tests
Target:	ABCA4
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of ATP Binding Cassette Transporter A4 (ABCA4). No significant cross-reactivity or interference between ATP Binding Cassette Transporter A4 (ABCA4) and analogues was observed.
Sensitivity:	0.123 ng/mL

Target Details

Target:	ABCA4
Alternative Name:	ATP Binding Cassette Transporter A4 (ABCA4 Products)

Target Details

larget Details		
Background:	Gene Name: ATP Binding Cassette Transporter A4 Gene Aliases: ABC-A4, ABC10, ABCR, ARMD2, CORD3, FFM, RMP, RP19, STGD, STGD1, Stargardt disease protein, Retinal-specific ATP-binding cassette transporter, RIM ABC transporter	
Gene ID:	24	
UniProt:	P78363	
Application Details		
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.	
Assay Time:	3 h	
Plate:	Pre-coated	
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to ATP Binding Cassette Transporter A4 (ABCA4). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to ATP Binding Cassette Transporter A4 (ABCA4). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplat well and incubated. After TMB substrate solution is added, only those wells that contain ATP Binding Cassette Transporter A4 (ABCA4), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically a a wavelength of 450nm ± 10nm. The concentration of ATP Binding Cassette Transporter A4 (ABCA4) in the samples is then determined by comparing the O.D. of the samples to the standard curve.	
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ATF Binding Cassette Transporter A4 (ABCA4) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ATP Binding Cassette Transporter A4 (ABCA4) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10%	

Application Details

	Inter-Assay: CV<12%	
Restrictions:	For Research Use only	
Handling		
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.	
Storage:	4 °C,-20 °C	
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.	
Expiry Date:	4-8 months	