

Datasheet for ABIN5652071

ABCC3 ELISA Kit



Go to Product page

()	11/	\sim	r\	ń	0	۱۸/	,
\cup	V	C	rv	ı	ヒ	٧V	

Quantity:	96 tests
Target:	ABCC3
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of ATP Binding Cassette Transporter C3 (ABCC3). No significant cross-reactivity or interference between ATP Binding Cassette Transporter C3 (ABCC3) and analogues was observed.
Sensitivity:	0.113 ng/mL

Target Details

Target:	ABCC3
Alternative Name:	ATP Binding Cassette Transporter C3 (ABCC3 Products)

Target Details

Background:	Gene Name: ATP Binding Cassette Transporter C3
	Gene Aliases: ABC31, MLP2, MOATD, MRP3, cMOAT2, Canalicular Multispecific Organic Anion
	Transporter 2, Multi-specific organic anion transporter D, Multidrug resistance-associated
	protein 3
Gene ID:	8714
UniProt:	015438
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5% within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to ATP Binding Cassette
	Transporter C3 (ABCC3). Standards or samples are then added to the appropriate microtiter
	plate wells with a biotin-conjugated antibody specific to ATP Binding Cassette Transporter C3
	(ABCC3). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate
	well and incubated. After TMB substrate solution is added, only those wells that contain ATP
	Binding Cassette Transporter C3 (ABCC3), biotin-conjugated antibody and enzyme-conjugated
	Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the
	addition of sulphuric acid solution and the color change is measured spectrophotometrically at
	a wavelength of 450nm ± 10nm. The concentration of ATP Binding Cassette Transporter C3
	(ABCC3) in the samples is then determined by comparing the O.D. of the samples to the
	standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ATF
	Binding Cassette Transporter C3 (ABCC3) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	ATP Binding Cassette Transporter C3 (ABCC3) were tested on 3 different plates, 8 replicates in
	each plate. CV(%) = SD/meanX100
	Intra-Assay: CV<10%

Application Details

	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months