

Datasheet for ABIN5652500 CAMKK2 ELISA Kit



Go to Product page

Overview

Quantity:	96 tests
Target:	CAMKK2
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Calcium/Calmodulin Dependent Protein Kinase Kinase 2 (CAMKK2). No significant cross-reactivity or interference between Calcium/Calmodulin Dependent Protein Kinase Kinase 2 (CAMKK2) and analogues was observed.
Sensitivity:	0.114 ng/mL

Target Details

Target:	CAMKK2
Alternative Name:	Calcium/Calmodulin Dependent Protein Kinase Kinase 2 (CAMKK2 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5652500 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Background:	Gene Name: Calcium/Calmodulin Dependent Protein Kinase Kinase 2
	Gene Aliases: CAMKK, CAMKKB, Calcium/calmodulin-dependent protein kinase kinase beta
Pathways:	AMPK Signaling
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, ai humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Calcium/Calmodulin Dependent Protein Kinase Kinase 2 (CAMKK2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Calcium/Calmodulin Dependent Protein Kinase Kinase 2 (CAMKK2). Next, Avidin conjugated to Calcium/Calmodulin Dependent Protein Kinase Kinase 2 (CAMKK2). Next, Avidin conjugated to the substrate solution is added, only those wells that contain Calcium/Calmodulin Dependent Protein Kinase Xinase 2 (CAMKK2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Calcium/Calmodulin Dependent Protein Kinase Xinase 2 (CAMKK2) in the samples is then determined by comparing the 0.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Calcium/Calmodulin Dependent Protein Kinase Kinase 2 (CAMKK2) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Calcium/Calmodulin Dependent Protein Kinase Kinase 2 (CAMKK2) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5652500 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Application Details	
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months