

Datasheet for ABIN5652591

CEACAM6 ELISA Kit



Overview

Quantity:	96 tests
Target:	CEACAM6
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	23.44 pg/mL - 1500 pg/mL
Minimum Detection Limit:	23.44 pg/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6). No significant cross-reactivity or interference between Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6) and analogues was observed.
Sensitivity:	9.68 pg/mL

Target Details

Target:	CEACAM6
Alternative Name:	Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6 Products)

Target Details Gene Name: Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 Background: Gene Aliases: CD66c, NCA, CD66-C, CEAL, Non-Specific Cross Reacting Antigen, Normal crossreacting antigen Gene ID: 4680 UniProt: P40199 **Application Details** Comment: The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end. Assay Time: 3 h Pre-coated Plate: Protocol: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate Related Cell Adhesion Molecule 6 (CEACAM6). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to

The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6) in the samples is then determined by comparing the O.D. of the samples to the standard curve.

Assay Precision:

Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6) were tested 20 times
on one plate, respectively
Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6) were tested on 3
different plates, 8 replicates in each plate. CV(%) = SD/meanX100

Application Details

	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months