

Datasheet for ABIN5652639

Casein Kinase 1 delta ELISA Kit



()	11/	\sim	r١.	/i	0	۱۸/	,
U	V	H	r٧	1	C	V۷	

Quantity:	96 tests
Target:	Casein Kinase 1 delta (CSNK1D)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Casein Kinase 1 Delta (CSNK1d). No significant cross-reactivity or interference between Casein Kinase 1 Delta (CSNK1d) and analogues was observed.
Sensitivity:	0.052 ng/mL

Target Details

Target:	Casein Kinase 1 delta (CSNK1D)
Alternative Name:	Casein Kinase 1 Delta (CSNK1D Products)

Target Details

Background:	Gene Name: Casein Kinase 1 Delta
	Gene Aliases: HCKID, Tau-protein kinase CSNK1D, Casein kinase I isoform delta
Gene ID:	1453
UniProt:	P48730
Pathways:	Hedgehog Signaling, M Phase
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Casein Kinase 1 Delta
	(CSNK1d). Standards or samples are then added to the appropriate microtiter plate wells with
	biotin-conjugated antibody specific to Casein Kinase 1 Delta (CSNK1d). Next, Avidin conjugated
	to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB
	substrate solution is added, only those wells that contain Casein Kinase 1 Delta (CSNK1d),
	biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The
	enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the colo
	change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The
	concentration of Casein Kinase 1 Delta (CSNK1d) in the samples is then determined by
	comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Casein Kinase 1 Delta (CSNK1d) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Casein Kinase 1 Delta (CSNK1d) were tested on 3 different plates, 8 replicates in each plate.
	CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only

Handling

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months