

Datasheet for ABIN5652871

CLIC1 ELISA Kit



Overview

Quantity:	96 tests
Target:	CLIC1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Chloride Intracellular Channel Protein 1 (CLIC1). No significant cross-reactivity or interference between Chloride Intracellular Channel Protein 1 (CLIC1) and analogues was observed.
Sensitivity:	0.116 ng/mL

Target Details

Target:	CLIC1
Alternative Name:	Chloride Intracellular Channel Protein 1 (CLIC1 Products)

Target Details

Background:	Gene Name: Chloride Intracellular Channel Protein 1
	Gene Aliases: G6, NCC27, p64CLCP, hRNCC, Chloride channel ABP, Nuclear chloride ion
	channel 27, Regulatory nuclear chloride ion channel protein
Gene ID:	1192
UniProt:	000299
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Chloride Intracellular
	Channel Protein 1 (CLIC1). Standards or samples are then added to the appropriate microtiter
	plate wells with a biotin-conjugated antibody specific to Chloride Intracellular Channel Protein 1
	(CLIC1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate
	well and incubated. After TMB substrate solution is added, only those wells that contain
	Chloride Intracellular Channel Protein 1 (CLIC1), biotin-conjugated antibody and enzyme-
	conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured
	spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Chloride
	Intracellular Channel Protein 1 (CLIC1) in the samples is then determined by comparing the O.D
	of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Chloride Intracellular Channel Protein 1 (CLIC1) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Chloride Intracellular Channel Protein 1 (CLIC1) were tested on 3 different plates, 8 replicates in
	each plate. CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%

Application Details

Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months