

Datasheet for ABIN5653205

C1QA ELISA Kit



Go to Product page

_				
	۱۱ / ۱	rv		۱۸/
	' V '	 ı v	Ι.	v v

Quantity:	96 tests
Target:	C1QA
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Complement Component 1, Q Subcomponent A (C1qA). No significant cross-reactivity or interference between Complement Component 1, Q Subcomponent A (C1qA) and analogues was observed.
Sensitivity:	0.059 ng/mL

Target Details

Target:	C1QA
Alternative Name:	Complement Component 1, Q Subcomponent A (C1QA Products)

Target Details

Background:	Gene Name: Complement Component 1, Q Subcomponent A	
	Gene Aliases: Complement Component 1,q Subcomponent,Alpha Polypeptide	
Gene ID:	712	
UniProt:	P02745	
Pathways:	Complement System	
Application Details		
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than	
	5 % within the expiration date under appropriate storage condition. To minimize extra influence	
	on the performance, operation procedures and lab conditions, especially room temperature, air	
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that	
	the whole assay is performed by the same operator from the beginning to the end.	
Assay Time:	3 h	
Plate:	Pre-coated	
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate	
	provided in this kit has been pre-coated with an antibody specific to Complement Component 1	
	Q Subcomponent A (C1qA). Standards or samples are then added to the appropriate microtiter	
	plate wells with a biotin-conjugated antibody specific to Complement Component 1, Q	
	Subcomponent A (C1qA). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to	
	each microplate well and incubated. After TMB substrate solution is added, only those wells	
	that contain Complement Component 1, Q Subcomponent A (C1qA), biotin-conjugated antibody	
	and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is	
	terminated by the addition of sulphuric acid solution and the color change is measured	
	spectrophotometrically at a wavelength of 450nm \pm 10nm. The concentration of Complement	
	Component 1, Q Subcomponent A (C1qA) in the samples is then determined by comparing the	
	O.D. of the samples to the standard curve.	
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level	
	Complement Component 1, Q Subcomponent A (C1qA) were tested 20 times on one plate,	
	respectively	
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level	
	Complement Component 1, Q Subcomponent A (C1qA) were tested on 3 different plates, 8	
	replicates in each plate. CV(%) = SD/meanX100	
	Intra-Assay: CV<10%	

Application Details

	Inter-Assay: CV<12%	
Restrictions:	For Research Use only	
Handling		
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.	
Storage:	4 °C,-20 °C	
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.	
Expiry Date:	4-8 months	