

## Datasheet for ABIN5653276 **CFHR1 ELISA Kit**



Overview

Quantity:	96 tests
Target:	CFHR1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.625 ng/mL - 40 ng/mL
Minimum Detection Limit:	0.625 ng/mL
Application:	ELISA

## Product Details

Sample Type:	Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Complement Factor H Related Protein 1 (CFHR1). No significant cross-reactivity or interference between Complement Factor H Related Protein 1 (CFHR1) and analogues was observed.
Sensitivity:	0.263 ng/mL
Target Details	

Target:	CFHR1
Alternative Name:	Complement Factor H Related Protein 1 (CFHR1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5653276 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details		
Background:	Gene Name: Complement Factor H Related Protein 1 Gene Aliases: CFHL, CFHL1, CFHL1P, CFHR1P, FHR1, H36-1, H36-2, HFL1, HFL2, H factor-like protein 1	
Application Details		
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.	
Assay Time:	3 h	
Plate:	Pre-coated	
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Complement Factor H Related Protein 1 (CFHR1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Complement Factor H Related Protein 1 (CFHR1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Complement Factor H Related Protein 1 (CFHR1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Complement Factor H Related Protein 1 (CFHR1) in the samples is then determined by comparing the O.D. of the samples to the standard curve.	
Assay Precision:	<ul> <li>Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level</li> <li>Complement Factor H Related Protein 1 (CFHR1) were tested 20 times on one plate,</li> <li>respectively</li> <li>Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level</li> <li>Complement Factor H Related Protein 1 (CFHR1) were tested on 3 different plates, 8 replicates</li> <li>in each plate. CV(%) = SD/meanX100</li> <li>Intra-Assay: CV&lt;10%</li> <li>Inter-Assay: CV&lt;12%</li> </ul>	
Restrictions:	For Research Use only	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5653276 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Н	land	lina
11	ana	mig

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months