

Datasheet for ABIN5653388

CRYBA1 ELISA Kit



Overview

Quantity:	96 tests
Target:	CRYBA1
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Crystallin Beta A1 (CRYbA1). No significant cross-reactivity or interference between Crystallin Beta A1 (CRYbA1) and analogues was observed.
Sensitivity:	0.118 ng/mL

Target Details

Target:	CRYBA1
Alternative Name:	Crystallin Beta A1 (CRYBA1 Products)

Target Details

Target Details	
Background:	Gene Name: Crystallin Beta A1
	Gene Aliases: CRYB1, Eye Lens Structural Protein
UniProt:	P02525
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, ai
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Crystallin Beta A1
	(CRYbA1). Standards or samples are then added to the appropriate microtiter plate wells with
	biotin-conjugated antibody specific to Crystallin Beta A1 (CRYbA1). Next, Avidin conjugated to
	Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB
	substrate solution is added, only those wells that contain Crystallin Beta A1 (CRYbA1), biotin-
	conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme
	substrate reaction is terminated by the addition of sulphuric acid solution and the color chang
	is measured spectrophotometrically at a wavelength of 450nm \pm 10nm. The concentration of
	Crystallin Beta A1 (CRYbA1) in the samples is then determined by comparing the O.D. of the
	samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Crystallin Beta A1 (CRYbA1) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Crystallin Beta A1 (CRYbA1) were tested on 3 different plates, 8 replicates in each plate. CV(%)
	= SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only

Handling

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months