

Datasheet for ABIN5653394 CUZD1 ELISA Kit



Overview

Quantity:	96 tests
Target:	CUZD1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Alternative Name:

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of CUB And Zona Pellucida Like Domains Protein 1 (CUZD1). No significant cross-reactivity or interference between CUB And Zona Pellucida Like Domains Protein 1 (CUZD1) and analogues was observed.
Sensitivity:	0.114 ng/mL
Target Details	
Target:	CUZD1

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5653394 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

CUB And Zona Pellucida Like Domains Protein 1 (CUZD1 Products)

Target Details			
Background:	Gene Name: CUB And Zona Pellucida Like Domains Protein 1		
	Gene Aliases: ERG-1, UO-44, Transmembrane protein UO-44		
Gene ID:	50624		
UniProt:	Q86UP6		
Application Details			
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than		
	5 % within the expiration date under appropriate storage condition. To minimize extra influence		
	on the performance, operation procedures and lab conditions, especially room temperature, air		
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that		
	the whole assay is performed by the same operator from the beginning to the end.		
Assay Time:	3 h		
Plate:	Pre-coated		
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate		
	provided in this kit has been pre-coated with an antibody specific to CUB And Zona Pellucida		
	Like Domains Protein 1 (CUZD1). Standards or samples are then added to the appropriate		
	microtiter plate wells with a biotin-conjugated antibody specific to CUB And Zona Pellucida Like		
	Domains Protein 1 (CUZD1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added		
	to each microplate well and incubated. After TMB substrate solution is added, only those wells		
	that contain CUB And Zona Pellucida Like Domains Protein 1 (CUZD1), biotin-conjugated		
	antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate		
	reaction is terminated by the addition of sulphuric acid solution and the color change is		
	measured spectrophotometrically at a wavelength of 450nm \pm 10nm. The concentration of		
	CUB And Zona Pellucida Like Domains Protein 1 (CUZD1) in the samples is then determined by		
	comparing the O.D. of the samples to the standard curve.		
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level		
	CUB And Zona Pellucida Like Domains Protein 1 (CUZD1) were tested 20 times on one plate,		
	respectively		
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level		
	CUB And Zona Pellucida Like Domains Protein 1 (CUZD1) were tested on 3 different plates, 8		
	replicates in each plate. CV(%) = SD/meanX100		
	Intra-Assay: CV<10%		
	Inter-Assay: CV<12%		

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5653394 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Application Details		
Restrictions:	For Research Use only	
Handling		
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.	
Storage:	4 °C,-20 °C	
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.	
Expiry Date:	4-8 months	