

Datasheet for ABIN5653462

CSRP1 ELISA Kit



Oo to rioduct page

Overviev	

Quantity:	96 tests
Target:	CSRP1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Cysteine And Glycine Rich Protein 1 (CSRP1). No significant cross-reactivity or interference between Cysteine And Glycine Rich Protein 1 (CSRP1) and analogues was observed.
Sensitivity:	0.057 ng/mL

Target Details

Target:	CSRP1
Alternative Name:	Cysteine And Glycine Rich Protein 1 (CSRP1 Products)

Target Details

Target Details	
Background:	Gene Name: Cysteine And Glycine Rich Protein 1 Cone Alianae: CRR1, CRR, CYRR, LIEL 141, Cysteine rich protein 1, Enididymia lyminal
	Gene Aliases: CRP1, CRP, CSRP, CYRP, HEL-141, Cysteine-rich protein 1, Epididymis luminal protein 141
Gene ID:	1465
UniProt:	P21291
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Cysteine And Glycine Rich
	Protein 1 (CSRP1). Standards or samples are then added to the appropriate microtiter plate
	wells with a biotin-conjugated antibody specific to Cysteine And Glycine Rich Protein 1 (CSRP1)
	Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and
	incubated. After TMB substrate solution is added, only those wells that contain Cysteine And
	Glycine Rich Protein 1 (CSRP1), biotin-conjugated antibody and enzyme-conjugated Avidin will
	exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of
	sulphuric acid solution and the color change is measured spectrophotometrically at a
	wavelength of 450nm \pm 10nm. The concentration of Cysteine And Glycine Rich Protein 1
	(CSRP1) in the samples is then determined by comparing the O.D. of the samples to the
	standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Cysteine And Glycine Rich Protein 1 (CSRP1) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Cysteine And Glycine Rich Protein 1 (CSRP1) were tested on 3 different plates, 8 replicates in
	each plate. CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%

Application Details

Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months